

cells proliferating to p68-86). Addition of recombinant IL-4 led to a significant decrease in IFN- $\gamma$  which was still dependent upon antigenic stimulation (Figures 4A and 4C  $0.19 \pm 0.08$  ng/ml without addition of MBP 68-86 versus  $2.37 \pm 0.8$  in cells proliferating to p68-86, a 12 fold increase).

5 Spleen T cells from anti-IGIF treated rats produced markedly reduced levels of IFN- $\gamma$  in response to antigenic stimulation in cultures that were or were not supplemented with IL-4 (Figure 4A,  $4.7 \pm 0.4$  ng/ml in spleen cells from anti-IGIF treated rat versus  $9.7 \pm 0.8$  in spleen cells from rats treated with normal rabbit IgG and  $13.5 \pm 0.7$  in PBS treated rats, with backgrounds of 10  $0.4$ ,  $0.8$  and  $0.7$ ,  $p < 0.001$ , when comparing anti-IGIF treatment to each control group). IL-4 production, however, markedly increased in splenic T cells from anti-IGIF treated rats regardless of *in vitro* stimulation (Figure 4B,  $62.3 \pm 4.2$  pg/ml in spleen cells from anti-IGIF treated rat versus  $15.3 \pm 0.4$  in spleen cells from rats treated with normal rabbit IgG and  $15.6 \pm 0.6$  in 15 PBS treated rats,  $p < 0.001$ , when comparing anti-IGIF treatment to each control group) unless cultures were supplemented with IL-4 (Figure 4D,  $1860 \pm 120$  pg/ml in spleen cells from anti-IGIF treated rat versus  $570 \pm 30$  in spleen cells from rats treated with normal rabbit IgG and  $450 \pm 35$  in PBS treated rats, with backgrounds of  $85$ ,  $42$  and  $34$ ,  $p < 0.0001$ , when 20 comparing anti-IGIF treatment to each control group). Addition of IL-4 to cultured spleen T cells (Figure 4C-D) did not exhibit a notable effect on their antigen specific proliferative response (data not shown).

TNF- $\alpha$  production was then evaluated in spleen cells from the above groups. The above spleen cells from anti-IGIF treated rats produced 25 markedly reduced levels of TNF- $\alpha$  in response to antigenic stimulation (Figure 5,  $850 \pm 45$  pg/ml in spleen cells from anti-IGIF treated rats versus  $1975 \pm 80$  in spleen cells from rats treated with normal rabbit IgG and  $2100 \pm 110$  in PBS treated rats, with backgrounds of  $230$ ,  $210$  and  $270$ , respectively,  $p < 0.001$ , when comparing anti-IGIF treatment to each control group). Thus perturbation of the Th1/Th2 balance in anti-IGIF treated rats 30 is associated with a marked reduction in TNF- $\alpha$  production.

Finally, the proliferative response of each group of cultured spleen cells to p68-86 was evaluated in a proliferation assay. ( $SI = 4.2 \pm 0.3$ ,  $3.6 \pm 0.4$  and  $3.96 \pm 0.5$  in spleen cells from rats treated with either anti-IGIF, 35 normal rabbit IgG or PBS respectively). Thus anti-IGIF immunotherapy alters Th2/Th1 balance without a notable affect on antigen specific proliferative responsiveness.

Although the invention has been described in conjunction with specific embodiments thereof, it is evident that many alternatives, modifications and variations will be apparent to those skilled in the art. Accordingly, it is intended to embrace all such alternatives, modifications and variations that fall within the spirit and broad scope of the appended claims.

5

0956798.002402

## REFERENCES CYTED

1. O'Garra, A., and K. Murphy. 1994. Role of cytokines in determining T-lymphocyte function. *Curr Opin Immunol* 6:458-66.
2. Abbas, A. K., K. M. Murphy, and A. Sher. 1996. Functional diversity of helper T lymphocytes. *Nature* 383:787-93.
3. Swain, S. L., A. D. Weinberg, M. English, and G. Huston. 1990. IL-4 directs the development of Th2-like helper effectors. *J. Immunol.* 145:3796-3806.
4. Lederer, J. A., V. L. Perez, L. DesRoches, S. M. Kim, A. K. Abbas, and A. H. Lichtman. 1996. Cytokine transcriptional events during helper T cell subset differentiation. *J Exp Med* 184:397-406.
5. Seder, R. A., R. Gazzinelli, A. Sher, and W. E. Paul. 1993. IL-12 acts directly on CD4<sup>+</sup> T cells to enhance priming for IFN- $\gamma$  production and diminishes IL-4 inhibition of such priming. *Proc. Natl. Acad. Sci. USA.* 90:10188-10192.
6. Seder, R. A., W. E. Paul, M. M. Davis, and B. Fazekas de St. Groth. 1992. The presence of interleukin-4 during in vitro priming determines the cytokine-producing potential of CD4<sup>+</sup> T cells from T cell receptor transgenic mice. *J. Exp. Med.* 176:1091-1098.
7. Fiorentino, D. F., A. Zlotnik, P. Vieira, T. R. Mosmann, M. Howard, K. W. Moore, and A. O'garra. 1991. IL-10 acts on the antigen presenting cell to inhibit cytokine production by Th1 cells. *J. Immunol.* 146:3444-3451.
8. Mosmann, T. R., and R. L. Coffman. 1989. Th1 and Th2 cells: different patterns of lymphokine secretion lead to different functional properties. *Annu.Rev.Immunol.* 9:145-173.
9. Mosmann, T., and K. Moor. 1989. The role of IL-10 in the crossregulation of Th1 and Th2 responses. *Immunol. Today.* 12:A49-A53.
10. Huang, H., J. Hu-Li, H. Chen, S. Z. Ben-Sasson, and W. E. Paul. 1997. IL-4 and IL-13 production in differentiated T helper type 2 cells is not IL-4 dependent. *J Immunol* 159:3731-8.
11. Fukaura, H., S. C. Kent, M. J. Pietrusiewicz, S. J. Khoury, H. L. Weiner, and D. A. Hafler. 1996. Induction of circulating myelin basic protein and proteolipid protein-specific transforming growth factor-beta1-secreting Th3 T cells by oral administration of myelin in multiple sclerosis patients. *Journal of Clinical Investigation* 98:70-7.